

DRUG FACTS

Active ingredient(s)

Purpose

Use(s)
treats head lice

Warnings
For external use only

Do not use
Do not use

- on children under 2 months of age
- near the eyes
- inside the nose, ear, mouth, or vagina
- on lice in eyebrows or eyelashes. See your doctor.

Ask a doctor before use if you have
Ask a doctor before use if you are

- allergic to ragweed. May cause breathing difficulty or an asthmatic episode.

Ask a doctor or pharmacist before use if you are

When using this product
When using this product

- keep eyes tightly closed and protect eyes with a washcloth or towel
- if product gets into the eyes, immediately flush with large amounts of water
- scalp itching or redness may occur

Stop use and ask a doctor if
Stop use and see a doctor if

- breathing difficulty occurs
- eye irritation occurs
- skin or scalp irritation continues or infection occurs

If pregnant or breast-feeding

Keep out of reach of children
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

Inspect

- all household members should be checked by another person for lice and/or nits (eggs)
- use a magnifying glass in bright light to help you see the lice and nits (eggs)
- use a tool, such as a comb or two unsharpened pencils to lift and part the hair
- look for tiny nits near the scalp, beginning at the back of the neck and behind the ears
- small sections of hair (1-2 inches wide) should be examined at a time
- unlike dandruff, nits stick to the hair. Dandruff should move when lightly touched.
- if either lice or nits (eggs) are found, treat with Nix Creme Rinse

Treat

- wash hair with a shampoo without conditioner. Do not use a shampoo that contains a conditioner or a conditioner alone since this may decrease the activity of Nix. Rinse with water.
- towel dry hair so it is damp but not wet
- shake the bottle of Nix well
- completely saturate the hair and scalp with Nix. Begin to apply Nix behind the ears and at the back of the neck.
- keep Nix out of the eyes. Protect the eyes with a washcloth or towel.
- leave Nix on the hair for 10 minutes, but no longer
- rinse with warm water
- towel dry hair and comb out tangles
- if live lice are seen seven days or more after the first treatment, a second treatment should be given

Remove Lice/Nits

- remove nits by combing the hair with the special small tooth comb provided. Remaining nits may be removed by hand (using a throw-away glove), or cutting the nits out.
- use the nit comb provided and make sure the hair remains slightly damp while removing nits
- if the hair dries during combing, dampen it slightly with water
- part the hair into 4 sections. Work on one section at a time. Longer hair may take more time (1-2 hours).
- start at the top of the head on the section you have picked
- with one hand, lift a 1-2 inch wide strand of hair. Get the teeth of the comb as close to the scalp as possible and comb with a firm, even motion away from the scalp to the end of the hair.
- use clips to pin back each strand of hair after you have combed out the nits
- clean the comb completely as you go. Wipe the nits from the comb with a tissue and throw away the tissue in a sealed plastic bag to prevent the lice from coming back.
- after combing, recheck the entire head for nits and repeat combing if necessary
- check the affected head daily to remove any nits that you might have missed

Other information

- read all the directions in the Consumer Information Insert and warnings before use. Keep the carton. It contains important information.
- store at 20° to 25°C (68° to 77°F)

Drug Facts

LIMITED CONSUMER WARRANTY

If you are not satisfied with Nix, we will refund your purchase price. Send cash register receipt, original UPC code, and reason for dissatisfaction to:

INSIGHT Pharmaceuticals Corp.
1170 Wheeler Way, Suite 150
Langhorne, PA 19047-1749

Distributed by:
INSIGHT Pharmaceuticals Corp.
Langhorne, PA 19047-1749
For a description of all of our products visit us at www.insightpharma.com

Inactive ingredients

balsam canada, cetyl alcohol, citric acid, FD&C yellow no. 6, fragrance, hydrolyzed animal protein, hydroxyethylcellulose, polyoxyethylene 10 cetyl ether, propylene glycol, stearalkonium chloride, water, isopropyl alcohol 5.6 g (20%), methylparaben 56 mg (0.2%), and propylparaben 22 mg (0.08%)